

Drinfel'd algebra deformations and the associahedra¹

Martin Markl and Steve Shnider

Introduction. The aim of this work is to construct a cohomology theory controlling the deformations of a general Drinfel'd algebra A and thus finish the program which began in [4], [5]. The picture presented here has two sides – the combinatorial one related with the fact of the existence of a differential graded Lie algebra structure on the simplicial cochain complex of the associahedra, and the algebraic one related with the algebra of derivations on the bar construction. The relationship between these two parts is described in Theorem 2.3. A full exposition of the results announced here will be given in [3].

1. Algebraic preliminaries.

Recall that a Drinfel'd algebra (or a quasi-bialgebra in the original terminology of [1]) is an object $A = (V, \cdot, \Delta, \Phi)$, where $(V, \cdot, \Delta)$ is an associative, not necessarily coassociative, unitary $\mathbf{k}$ -bialgebra, Φ is an invertible element of $V^{\otimes 3}$, and the usual coassociativity property is replaced by

$$(\mathbb{1} \otimes \Delta)\Delta \cdot \Phi = \Phi \cdot (\Delta \otimes \mathbb{1})\Delta,$$

where we use the dot $\cdot$ to indicate both the (associative) multiplication on V and the induced multiplication on $V^{\otimes 3}$. Moreover, the validity of the following “pentagon identity” is required:

$$(\mathbb{1}^2 \otimes \Delta)(\Phi) \cdot (\Delta \otimes \mathbb{1}^2)(\Phi) = (1 \otimes \Phi) \cdot (\mathbb{1} \otimes \Delta \otimes \mathbb{1})(\Phi) \cdot (\Phi \otimes 1).$$

For a $(V, \cdot)$ -module N , define the following generalization of the M -construction of [4, par. 3]. Let $F^* = \bigoplus_{n \geq 0} F^n$ be the free unitary nonassociative $\mathbf{k}$ -algebra generated by N , graded by length of words. The algebra F^* admits a natural left action, $(a, f) \mapsto a \bullet f$, of $(V, \cdot)$ given by the rules:

- (i) on $F^1 = N$, the action is given by the action of V on N and
- (ii) $a \bullet (f \star g) = \sum (\Delta'(a) \bullet f) \star (\Delta''(a) \bullet g)$,

where $\star$ stands for the multiplication in F^* and we use the notation $\Delta(a) = \sum \Delta'(a) \otimes \Delta''(a)$. The right action $(f, b) \mapsto f \bullet b$ is defined by the similar rules. These actions define on F^* the structure of a $(V, \cdot)$ -bimodule.

Let $\sim$ be the relation, $\star$ -multiplicatively generated on F^* by the expressions of the form

$$(1) \quad \left((\Phi_1 \bullet x) \star ((\Phi_2 \bullet y) \star (\Phi_3 \bullet z)) \right) \sim \left(((x \bullet \Phi_1) \star (y \bullet \Phi_2)) \star (z \bullet \Phi_3) \right),$$

where $\Phi = \sum \Phi_1 \otimes \Phi_2 \otimes \Phi_3$ and $x, y, z \in F^*$. Put $\odot(N) := F^* / \sim$. Just as in [4, Proposition 3.2] one proves that the $\bullet$ -action induces on $\odot(N)$ the structure of a $(V, \cdot)$ -bimodule (denoted again by $\bullet$) and

¹First author partially supported by the E. Schrödinger International Institute for Mathematical Physics, Vienna, and by the National Research Counsel, USA

that $\star$ induces on $\odot(N)$ the nonassociative multiplication denoted by $\odot$. The operations are related by

$$(2) \quad a \bullet (f \odot g) = (\Delta'(a) \bullet f) \odot (\Delta''(a) \bullet g) \text{ and } (f \odot g) \bullet b = (f \bullet \Delta'(b)) \odot (g \bullet \Delta''(b)),$$

for $a, b \in V$ and $f, g \in \odot(N)$. Since the defining relations (1) are homogeneous with respect to the length, the grading of F^* induces on $\odot(N)$ the grading $\odot^*(N) = \bigoplus_{i \geq 0} \odot^i(N)$. If N itself is a graded vector space, we have also the obvious second grading, $\odot(N) = \bigoplus_j \odot(N)^j$, which coincides with the first grading if N is concentrated in degree 1.

Let $\text{Der}_V^n(\odot N)$ denote the set of degree n $(V, \cdot)$ -linear derivations of the (nonassociative) graded algebra $\odot(N)^*$. It is immediate to see that there is an one-to-one correspondence between the elements $\theta \in \text{Der}_V^n(\odot N)$ and $(V, \cdot)$ -linear homogeneous degree n maps $f : N^* \rightarrow \odot(N)^*$.

If $N = X \oplus Y$, then $\odot(X \oplus Y)$ is naturally bigraded, $\odot^{*,*}(X \oplus Y) = \bigoplus_{i,j \geq 0} \odot^{i,j}(X \oplus Y)$, this bigrading being defined by saying that a monomial w belongs to $\odot^{i,j}(X \oplus Y)$ if there are exactly i (resp. j) occurrences of the elements of X (resp. Y) in w .

Let $(\mathcal{B}(V), d_{\mathcal{B}})$ be the two-sided bar construction on the algebra $(V, \cdot)$, i.e. the graded differential space $\mathcal{B}(V) = \bigoplus_{n \geq 1} \mathcal{B}_n(V)$, where $\mathcal{B}_n(V)$ is the free $(V, \cdot)$ -bimodule on $V^{\otimes n}$, i.e. the vector space $V^{\otimes(n+2)}$ with the action given by

$$u \cdot (a_0 \otimes \cdots \otimes a_{n+1}) := (u \cdot a_0 \otimes \cdots \otimes a_{n+1}) \text{ and } (a_0 \otimes \cdots \otimes a_{n+1}) \cdot w := (a_0 \otimes \cdots \otimes a_{n+1} \cdot w)$$

for $u, v, a_0, \dots, a_{n+1} \in V$. If we use the more compact notation (though a nonstandard one), writing $(a_0 | \cdots | a_{n+1})$ instead of $a_0 \otimes \cdots \otimes a_{n+1}$, the differential $d_{\mathcal{B}} : \mathcal{B}_n(V) \rightarrow \mathcal{B}_{n-1}(V)$ is defined as

$$d_{\mathcal{B}}(a_0 | \cdots | a_{n+1}) := \sum_{0 \leq i \leq n} (-1)^i (a_0 | \cdots | a_i \cdot a_{i+1} | \cdots | a_{n+1}).$$

Notice that the differential $d_{\mathcal{B}}$ is a $(V, \cdot)$ -linear map. We have two more $(V, \cdot)$ -linear maps, namely the ‘coactions’ $\lambda : \mathcal{B}(V) \rightarrow V \odot \mathcal{B}(V)$ and $\rho : \mathcal{B}(V) \rightarrow \mathcal{B}(V) \odot V$ given by

$$\begin{aligned} \lambda(a_0 | \cdots | a_{n+1}) &:= \sum \Delta'(a_0) \cdots \Delta'(a_{n+1}) \odot (\Delta''(a_0) | \cdots | \Delta''(a_{n+1})), \text{ and} \\ \rho(a_0 | \cdots | a_{n+1}) &:= \sum (\Delta'(a_0) | \cdots | \Delta'(a_{n+1})) \odot \Delta''(a_0) \cdots \Delta''(a_{n+1}). \end{aligned}$$

The following notation will be useful in the sequel. Let $\uparrow V$ denotes the vector space V considered as a graded vector space concentrated in degree 1 and let $\tilde{\mathcal{B}}(V)$ be the bar construction $\mathcal{B}(V)$ with the ‘opposite’ grading, $(\tilde{\mathcal{B}}(V))^i = \mathcal{B}_{-i}(V)$. Then put $\odot(V, \mathcal{B}(V)) := \odot(\uparrow V \oplus \tilde{\mathcal{B}}(V))$. The maps above determine the following two important elements of $\text{Der}_V^1(\odot(V, \mathcal{B}(V)))$: the derivation D_{-1} given by $D_{-1}|_{\mathcal{B}(V)} = d_{\mathcal{B}}$ and $D_{-1}|_V = 0$, and the derivation D_0 given by $D_0|_{\mathcal{B}_n(V)} = \lambda + (-1)^{n+1}\rho$, $1 \leq n$, and $D_0|_V = \Delta$.

Theorem 1.1. *There are derivations $D_k \in \text{Der}_V^1(\odot(V, \mathcal{B}(V)))$, $k \geq 1$, such that $D_k(\odot^{i,j}(V, \mathcal{B}_n(V))) \subset \odot^{i+k+1,j}(V, \mathcal{B}_{n+k}(V))$ for $i, j \geq 0$, $k, n \geq 1$, and that $D := D_{-1} + D_0 + \sum_{k \geq 1} D_k$ has square zero, $D^2 = 0$.*

The construction of D_k is described in the next section.

2. On the Stasheff associahedra and the higher derivations D_k .

Let K_n be, for $n \geq 2$, the Stasheff associahedron [6]. The vertices of K_n are indexed by the set Br_n of full bracketings of n indeterminates. For $a, b \geq 2$ and $1 \leq t \leq b$, there is a map $(-, -)_t : \text{Br}_a \times \text{Br}_b \rightarrow \text{Br}_{a+b-1}$, given by inserting the first argument to the t -th position in the second argument. This map defines the inclusions $\iota_t : K_a \times K_b \hookrightarrow \partial K_{a+b-1}$ and it is known that

$$(3) \quad \partial K_n = \bigcup_J \iota_t(K_a \times K_b),$$

where $J = \{(a, b, t); a + b = n + 1, a, b \geq 2 \text{ and } 1 \leq t \leq b\}$.

Let $\prec$ be the partial order on Br_n defined by saying that $u \prec v$ if and only if v is obtained from u by the substitution $(w_1, w_2)w_3 \mapsto w_1(w_2, w_3)$ with some $w_i \in \text{Br}_{n_i}$, $i = 1, 2, 3$, $n_1 + n_2 + n_3 \leq n$. Let ξ_n be the (unique) minimal element of Br_n .

Proposition 2.1.

$$K_n = \text{Cone}(\bigcup_{J'} \iota_t(K_a \times K_b); \xi_n),$$

where $J' = \{(a, b, t) \in J; t \geq 2\}$.

The cone structure above induces on K_n inductively the triangulation $\mathcal{T}_n$ as follows:

- (i) $\mathcal{T}_2$ is the only possible triangulation of $K_2 = \{\bullet\bullet\}$ and
- (ii) for $n \geq 2$, $\mathcal{T}_n$ is induced by the cone structure of Proposition 2.1 from the triangulations $\mathcal{T}_a \times \mathcal{T}_b$ of $K_a \times K_b$, $a + b = n + 1$, $a, b \geq 2$, which are already defined, by the induction assumption.

It is possible to show that $\mathcal{T}_n$ has the property that $\prec$ induces a total order on the vertices of any simplex of $\mathcal{T}_n$. We will consider $\mathcal{T}_n$ as an oriented triangulation with the orientation induced by $\prec$.

Let $C_i(K_n)$ denote the set of i -dimensional oriented simplicial chains of $\mathcal{T}_n$ with coefficients in $\mathbf{k}$ and let $d_S : C_i(K_n) \rightarrow C_{i-1}(K_n)$ be the simplicial differential. For $\mathbf{s} \in C_p(K_a)$ and $\mathbf{t} \in C_q(K_b)$, $p, q \geq 0$, $a, b \geq 2$, set

$$\mathbf{s} \diamond \mathbf{t} := \sum_{1 \leq t \leq b} (-1)^{(a+1)(t+q+1)} (\iota_t)_* (\mathbf{s} \times \mathbf{t}) \in C_{p+q}(K_{a+b-1}).$$

Let $C(K)^n := \bigoplus_{n=i-p-1} C_p(K_i)$ and extend $\diamond$ to a bilinear map (denoted by the same symbol) $\diamond : C(K)^* \otimes C(K)^* \rightarrow C(K)^*$.

Proposition 2.2. $(C(K)^*, \diamond, d_S)$ is a graded differential pre-Lie algebra (an analog of the corresponding nondifferential notion of [2]), i.e.

$$\begin{aligned} \mathbf{s} \diamond (\mathbf{t} \diamond \mathbf{u}) - (\mathbf{s} \diamond \mathbf{t}) \diamond \mathbf{u} &= (-1)^{|\mathbf{s}| \cdot |\mathbf{t}|} (\mathbf{t} \diamond (\mathbf{s} \diamond \mathbf{u}) - (\mathbf{t} \diamond \mathbf{s}) \diamond \mathbf{u}) \text{ and} \\ d_S(\mathbf{s} \diamond \mathbf{t}) &= d_S \mathbf{s} \diamond \mathbf{t} + (-1)^{|\mathbf{s}|} \cdot \mathbf{s} \diamond d_S \mathbf{t} \end{aligned}$$

for any $\mathbf{s}, \mathbf{t}, \mathbf{u} \in C(K)^*$.

Let $[\mathbf{s}, \mathbf{t}] := \mathbf{s} \diamond \mathbf{t} - (-1)^{|\mathbf{s}| \cdot |\mathbf{t}|} \mathbf{t} \diamond \mathbf{s}$. An easy computation based on Proposition 2.2 shows that $(C(K)^*, [-, -], d_S)$ is a differential graded Lie algebra. Observe that $\text{Der}_V^*(\odot(V, \mathcal{B}(V)))$ is also a differential graded Lie algebra, the bracket being given by the (graded) commutator of derivations and the differential defined by $\nabla(\theta) := [D_{-1}, \theta]$.

Let $\mathbf{e}_n \in C_n(K_{n+2}) \subset C(K)^1$, $n \geq 0$, be a sequence of ‘fundamental classes’ chosen in such a way that $\mathbf{e}_0 = 1 \cdot (\bullet, \bullet)$ and that

$$(4) \quad d_S \mathbf{e}_n + \sum_{i+j=n-1} (\mathbf{e}_i \diamond \mathbf{e}_j) = 0, \text{ or, equivalently, } d_S \mathbf{e}_n + \frac{1}{2} \sum_{i+j=n-1} [\mathbf{e}_i, \mathbf{e}_j].$$

It follows from (3) and from the asphericity of K_n that such a sequence exists.

Theorem 2.3. *There exists a homomorphism $m: (C(K)^*, [-, -], d_S) \rightarrow (\text{Der}_V^*(\odot(V, \mathcal{B}(V))), [-, -], \nabla)$ of differential graded Lie algebras such that $m(\mathbf{e}_0) = D_0$. Moreover, the homomorphism m has the property that*

$$m(C_p(K_n))(\odot^{i,j}(V, \mathcal{B}_q(V))) \subset \odot^{i+n-1,j}(V, \mathcal{B}_{p+q}(V)), \quad i, j, p \geq 0, \quad n \geq 2, \quad q \geq 1.$$

Theorem 2.4. *The derivations $D_k := m(\mathbf{e}_k)$, $k \geq 1$, satisfy the assumptions of Theorem 1.1.*

Sketch of proof. Looking at the homogeneous components of D^2 we see that $D^2 = 0$ is equivalent to $[D_{-1}, D_k] + \frac{1}{2} \sum_{i+j=k-1} [D_i, D_j] = 0$ which is, by the definition of the derivations D_k and Theorem 2.3, the same as $m(d_S \mathbf{e}_k + \frac{1}{2} \sum_{i+j=k-1} [\mathbf{e}_i, \mathbf{e}_j]) = 0$, which follows from (4). ■

3. Deformation theory for Drinfel’d algebras.

As we have already observed, both $\odot(V, \mathcal{B}(V))^*$ and $\odot^*(V) (= \odot(\uparrow V)^*)$ are graded $(V, \cdot)$ -modules (via the $\bullet$ -action). The $\odot$ -multiplication yields on both $\odot(V, \mathcal{B}(V))^*$ and $\odot^*(V)$ the structures of graded $\odot^*(V)$ -bimodules and both structures are related by (2). Denote by $\mathbf{Hom}^n(\odot(V, \mathcal{B}(V))^*, \odot^*(V))$ the set of all degree n homogeneous maps $f: \odot(V, \mathcal{B}(V))^* \rightarrow \odot^*(V)$ which are both $\odot^*(V)$ and $(V, \cdot)$ -linear. For such a map put $d(f) := f \circ D + (-1)^n d_C \circ f$, where D is as in Theorem 1.1 and d_C is the degree one derivation on $\odot^*(V)$ defined by $d_C|_V := \Delta$.

Theorem 3.1. *The deformation theory of a Drinfel’d algebra $A = (V, \cdot, \Delta, \Phi)$ is controlled by the cohomology of the complex*

$$(\mathbf{Hom}^*(\odot(V, \mathcal{B}(V))^*, \odot^*(V)), d).$$

Using the fact that $\odot(V, \mathcal{B}(V))^*$ is, in a sense, the free $\odot^*(V)$ -bimodule on the $(V, \cdot)$ -bimodule $\tilde{\mathcal{B}}(V)^*$ we may write

$$\mathbf{Hom}^n(\odot(V, \mathcal{B}(V))^*, \odot^*(V)) \cong \text{Hom}_V^n(\tilde{\mathcal{B}}(V)^*, \odot^*(V)),$$

where $\text{Hom}_V^n(\tilde{\mathcal{B}}(V)^*, \odot^*(V))$ denotes the set of $(V, \cdot)$ -linear maps $f: \tilde{\mathcal{B}}(V) \rightarrow \odot^*(V)$. Using the fact that $\tilde{\mathcal{B}}(V)^*$ is the free $(V, \cdot)$ -module on $\otimes^{-*}(V)$ we get the identification

$$\text{Hom}_V^n(\tilde{\mathcal{B}}(V)^*, \odot^*(V)) \cong \text{Hom}_k^n(\otimes^{-*}(V), \odot^*(V)),$$

where $\text{Hom}_{\mathbf{k}}^n(\otimes^{-*}(V), \odot^*(V))$ denotes the set of degree n $\mathbf{k}$ -linear homogeneous maps $g : \otimes^{-*}(V) \rightarrow \odot^*(V)$. Finally, using the canonical isomorphism $J : \odot^*(V) \cong \otimes^*(V)$ of graded vector spaces constructed in [4, Proposition 3.3] we get

$$\text{Hom}_{\mathbf{k}}^n(\otimes^{-*}(V), \odot^*(V)) \cong \text{Hom}_{\mathbf{k}}^n(\otimes^{-*}(V), \otimes^*(V)),$$

where we denote the corresponding differential again by d . This is the underlying vector space of the Gerstenhaber-Schack complex [7] which controls the deformation of coassociative bialgebras. In fact, our complex can be considered as a kind of deformation of the Gerstenhaber-Schack complex as follows. In order to understand better the structure of the differential graded space $(\text{Hom}_{\mathbf{k}}^n(\otimes^{-*}(V), \otimes^*(V)), d)$, let us simplify the notation by setting

$$C^n(A) := \text{Hom}_{\mathbf{k}}^n(\otimes^{-*}(V), \otimes^*(V)) \text{ and } E^{i,j}(A) := \text{Hom}_{\mathbf{k}}(\otimes^i(V), \otimes^j(V)).$$

Then it is immediate to see that $C^n(A) = \bigoplus_{i+j=n} E^{i,j}(A)$ and that the differential d decomposes as $d = d_{\text{Hoch}} + d_{\text{q-coH}} + \sum_{k=1}^{\infty} M^k$, where

$$(5) \quad \begin{aligned} f &\mapsto f \circ D_{-1} && \text{induces } d_{\text{Hoch}} : E^{i,j}(A) \rightarrow E^{i+1,j}(A), \\ f &\mapsto f \circ D_0 + (-1)^n d_C \circ f && \text{induces } d_{\text{q-coH}} : E^{i,j}(A) \rightarrow E^{i,j+1}(A) \\ &\text{and } f \mapsto f \circ D_k && \text{induces } M^k : E^{i,j}(A) \rightarrow E^{i-k,j+k+1}(A). \end{aligned}$$

Here d_{Hoch} is the Hochschild differential of the associative algebra $(V, \cdot)$, $d_{\text{q-coH}}$ is the analog of the coHochschild differential for the (noncoassociative) coalgebra (V, Δ) , and the maps M^k correspond to the derivations D_k of Theorem 1.1, for $k \geq 1$. In general, when $\Phi \neq 1$, we do not have $d_{\text{q-coH}}^2 = 0$. However when $\Phi = 1$, $d_{\text{q-coH}}^2 = 0$ and $d_{\text{Hoch}} + d_{\text{q-coH}}$ is the differential defined in [7]. Moreover, in this case the operators D_k vanish when restricted to normalized cochains, so the cohomology of our complex is the same as that of the Gerstenhaber-Schack complex.

Bigraded differential complexes where the differential decomposes in as in (5) are sometimes called hypercomplexes (although the first author prefers to call them monsters). Theorem 3.1 can be then reformulated as

Theorem 3.1.' *The deformation theory of a Drinfel'd algebra $A = (V, \cdot, \Delta, \Phi)$ is controlled by the cohomology of the hypercomplex*

$$(C^*(A) = \bigoplus_{i+j=*} E^{i,j}(A), d = d_{\text{Hoch}} + d_{\text{q-coH}} + \sum_{k=1}^{\infty} M^k)$$

References.

- [1] V.G. Drinfel'd. Kvazichopfovy algebr. *Algebra i Analiz*, **1,6**(1989), 114–148.
- [2] G. Gerstenhaber. The cohomology structure of an associative rings. *Annals of Mathematics*, **78,2**(1963), 268–288.
- [3] M. Markl and S. Shnider. Drinfel'd algebra deformations, homotopy comodules and the associahedra. To appear.

- [4] M. Markl and J.D. Stasheff. Deformation theory via deviations. *Journal of Algebra*, to appear.
- [5] S. Shnider and S. Sternberg. The cobar resolution and a restricted deformation theory for Drinfeld algebras. *Journal of Algebra*, to appear.
- [6] J.D. Stasheff. Homotopy associativity of H-spaces I,II. *Trans. Amer. Math. Soc.*, **108**(1963), 275–312.
- [7] M. Gerstenhaber and S. D. Schack. Bialgebra Cohomology, Deformations, and Quantum Groups. *Proc. Nat. Acad. Sci.*, **87**(1990), 478–481.

M. M.: Mathematical Institute of the Academy, Žitná 25, 115 67 Praha 1, Czech Republic,
email: **markl@earn.cvut.cz**

Current address (until March 1994): Math-UNC, Chapel Hill, NC 27599-3250, USA

S. S.: Department of Mathematics, University of Bar Ilan, Israel,
email: **shnider@bimacs.cs.biu.ac.il**